

兔抗 BCL2A1 多克隆抗体

中文名称：兔抗 BCL2A1 多克隆抗体

英文名称：Anti-BCL2A1 rabbit polyclonal antibody

别名：GRS; BFL1; ACC-1; ACC-2; HBPA1; BCL2L5

抗原：BCL2A1

储存：冷冻（-20℃）

宿主：Rabbit

相关类别：一抗

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	<p>This gene encodes a member of the BCL-2 protein family. The proteins of this family form hetero- or homodimers and act as anti- and pro-apoptotic regulators that are involved in a wide variety of cellular activities such as embryonic development, homeostasis and tumorigenesis. The protein encoded by this gene is able to reduce the release of pro-apoptotic cytochrome c from mitochondria and block caspase activation. This gene is a direct transcription target of NF-kappa B in response to inflammatory mediators, and is up-regulated by different extracellular signals, such as granulocyte-macrophage colony-stimulating factor (GM-CSF), CD40, phorbol ester and inflammatory cytokine TNF and IL-1, which suggests a cytoprotective function that is e</p>
--------------------	---

	essential for lymphocyte activation as well as cell survival. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.
Applications:	ELISA, WB, IHC
Name of antibody:	BCL2A1
Immunogen:	Fusion protein of human BCL2A1
Full name:	BCL2-related protein A1
Synonyms :	GRS; BFL1; ACC-1; ACC-2; HBPA1; BCL2L5
SwissProt:	Q16548
ELISA Recommended dilution:	2000-5000
IHC positive control:	Human thyroid cancer
IHC Recommend dilution:	25-100
WB Predicted band size:	20 kDa
WB Positive control:	Mouse testis tissue and hela cells
WB Recommended dilution:	200-1000



