

## 兔抗 DRG1 多克隆抗体

中文名称: 兔抗 DRG1 多克隆抗体

英文名称: Anti-DRG1 rabbit polyclonal antibody

别 名: NEDD3

相关类别: 一抗

抗原: DRG1

储 存: 冷冻(-20℃)

宿 主: Rabbit

反应种属: Human, Mouse

标记物: Unconjugate

克隆类型: rabbit polyclonal

## 技术规格

belongs to the GTP1/OBG family. Expressed at high lev els in heart, kidney and skeletal muscle and at lower le vels in brain, liver, placenta, lung, colon and spleen, D RG1 binds to TAL1 and TAL2 and is thought to play a role in cell proliferation and differentiation, as well as in apoptosis, suggesting a role in tumor formation and


metastasis. DRG1 is subject to polyubiquitination and s umoylation, the former of which induces proteolytic de gradation. The gene encoding DRG1 maps to human c

DRG1 (developmentally regulated GTP binding protein 1), also known as NEDD3 (neural precursor cell express ed developmentally down-regulated protein 3), is a 367 amino acid protein that localizes to the cytoplasm and



## 全国订货电话 4008-723-722

p	.,,
	hromosome 22, which houses over 500 genes and is the second smallest human chromosome. Mutations in several of the genes that map to chromosome 22 are involved in the development of Phelan-McDermid syndrome, Neurofibromatosis type 2, autism and schizophrenia.
Applications:	ELISA, WB, IHC
Name of antibody:	DRG1
Immunogen:	Fusion protein of human DRG1
Full name:	developmentally regulated GTP binding protein 1
Synonyms:	NEDD3
SwissProt:	Q9Y295
ELISA Recommended dilution:	2000-5000
IHC positive control:	Human thyroid cancer and human liver cancer
IHC Recommend dilution:	50-200
WB Predicted band size:	41 kDa
WB Positive control:	Human fetal liver and brain tissue, 293T and Hela cells
WB Recommended dilution:	500-2000





## 全国订货电话 4008-723-722

